

F(ab')₂ fragments

¹²³I 13.2 h

Organ	Absorbed dose per unit activity administered (mGy/MBq)				
	Adult	15 years	10 years	5 years	1 year
Adrenals	2.1E-02	2.7E-02	4.1E-02	6.1E-02	1.1E-01
Bladder	4.7E-02	6.1E-02	8.9E-02	1.3E-01	2.4E-01
Bone surfaces	1.9E-02	2.3E-02	3.6E-02	5.9E-02	1.1E-01
Brain	5.6E-03	7.0E-03	1.1E-02	1.9E-02	3.4E-02
Breasts	5.6E-03	7.0E-03	1.1E-02	1.7E-02	3.3E-02
Gallbladder	2.3E-02	2.8E-02	4.0E-02	6.0E-02	1.0E-01
Gastrointestinal tract					
Stomach	1.2E-02	1.6E-02	2.5E-02	3.8E-02	6.7E-02
Small intestine	1.1E-02	1.4E-02	2.3E-02	3.7E-02	6.5E-02
Colon	1.1E-02	1.4E-02	2.2E-02	3.5E-02	6.2E-02
(Upper large intestine)	1.2E-02	1.5E-02	2.4E-02	3.9E-02	6.8E-02
(Lower large intestine)	9.8E-03	1.2E-02	1.9E-02	3.0E-02	5.4E-02
Heart	1.1E-02	1.3E-02	2.1E-02	3.1E-02	5.7E-02
Kidneys	1.7E-01	2.0E-01	2.8E-01	4.1E-01	7.1E-01
Liver	6.0E-02	7.7E-02	1.2E-01	1.6E-01	2.9E-01
Lungs	1.0E-02	1.3E-02	2.0E-02	3.0E-02	5.6E-02
Muscles	8.3E-03	1.0E-02	1.6E-02	2.4E-02	4.6E-02
Oesophagus	7.0E-03	8.9E-03	1.4E-02	2.1E-02	3.9E-02
Ovaries	1.0E-02	1.3E-02	2.0E-02	3.2E-02	5.8E-02
Pancreas	2.0E-02	2.5E-02	3.9E-02	5.9E-02	1.0E-01
Red marrow	1.7E-02	1.9E-02	3.0E-02	5.2E-02	1.1E-01
Skin	5.0E-03	6.1E-03	9.7E-03	1.6E-02	3.0E-02
Spleen	9.3E-02	1.3E-01	2.0E-01	3.0E-01	5.3E-01
Testes	6.4E-03	8.3E-03	1.3E-02	2.1E-02	4.1E-02
Thymus	7.0E-03	8.9E-03	1.4E-02	2.1E-02	3.9E-02
Thyroid	6.4E-03	8.1E-03	1.3E-02	2.1E-02	4.0E-02
Uterus	1.2E-02	1.5E-02	2.4E-02	3.8E-02	6.8E-02
Remaining organs	8.8E-03	1.1E-02	1.7E-02	2.7E-02	4.8E-02
Effective dose (mSv/MBq)	1.9E-02	2.3E-02	3.5E-02	5.4E-02	9.9E-02